



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, 15/10, C07K 14/415	A2	(11) International Publication Number: WO 99/29882 (43) International Publication Date: 17 June 1999 (17.06.99)																																																																																																																																																																																																																																								
(21) International Application Number: PCT/US98/26209 (22) International Filing Date: 10 December 1998 (10.12.98) (30) Priority Data: 08/988,015 10 December 1997 (10.12.97) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/988,015 (CON) Filed on 10 December 1997 (10.12.97) (71) Applicant (for all designated States except US): PIONEER HI-BRED INTERNATIONAL, INC. [US/US]; 800 Capital Square, 400 Locust Street, Des Moines, IA 50309 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): RAO, Gururaj, A. [US/US]; 4734 74th Street, Urbandale, IA 50322 (US). SLEISTER, Heidi, Major [US/US]; 1302 N.W. Cedarwood Drive, Ankeny, IA 50021 (US). (74) Agents: SPRUILL, W., Murray et al.; Alston & Bird LLP, P.O. Drawer 34009, Charlotte, NC 28234-4009 (US).		(81) Designated States: AL, AM, AT, AT (Utility model), AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, CZ (Utility model), DE, DE (Utility model), DK, DK (Utility model), EE, EE (Utility model), ES, FI, FI (Utility model), GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SK (Utility model), SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>																																																																																																																																																																																																																																								
(54) Title: METHODS FOR ALTERING THE NUTRITIONAL VALUE OF A PLANT PROTEIN, VEGETATIVE STORAGE PROTEIN, VSP, BY ALTERING THE AMINO ACID CONTENT OF PROTEINS																																																																																																																																																																																																																																										
(57) Abstract <p>Methods and compositions for altering amino acid composition of a protein of interest are provided, particularly proteins whose three-dimensional structure is unknown. The method comprises creating interacting molecules to the native protein and selecting for engineered proteins which retain the native conformation by antibody binding. In this manner, the levels of essential amino acids in a protein can be increased yet the biological activity of the protein maintained.</p> <div data-bbox="792 1161 1352 1833"> <p style="text-align: center;">VSP HOMOLOGIES</p> <table border="1"> <thead> <tr> <th></th> <th>1</th> <th>5</th> <th>10</th> <th>15</th> <th>20</th> <th>25</th> <th>30</th> </tr> </thead> <tbody> <tr><td>VSP-b</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>VSP-a</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>T.phos</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ph.vulg</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ar.VSP</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ar.1A-1</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ar.17A-1</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> </tbody> </table> <table border="1"> <thead> <tr> <th></th> <th>31</th> <th>35</th> <th>40</th> <th>45</th> <th>50</th> <th>55</th> </tr> </thead> <tbody> <tr><td>VSP-b</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>VSP-a</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>T.phos</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ph.vulg</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ar.VSP</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ar.1A-1</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ar.17A-1</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> </tbody> </table> <table border="1"> <thead> <tr> <th></th> <th>60</th> <th>65</th> <th>70</th> <th>75</th> <th>80</th> <th>85</th> </tr> </thead> <tbody> <tr><td>VSP-b</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>VSP-a</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>T.phos</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ph.vulg</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ar.VSP</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ar.1A-1</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ar.17A-1</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> </tbody> </table> <table border="1"> <thead> <tr> <th></th> <th>90</th> <th>95</th> <th>100</th> <th>105</th> <th>110</th> <th>115</th> </tr> </thead> <tbody> <tr><td>VSP-b</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>VSP-a</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>T.phos</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ph.vulg</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ar.VSP</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ar.1A-1</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> <tr><td>Ar.17A-1</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td><td>S</td></tr> </tbody> </table> </div>				1	5	10	15	20	25	30	VSP-b	S	S	S	S	S	S	S	VSP-a	S	S	S	S	S	S	S	T.phos	S	S	S	S	S	S	S	Ph.vulg	S	S	S	S	S	S	S	Ar.VSP	S	S	S	S	S	S	S	Ar.1A-1	S	S	S	S	S	S	S	Ar.17A-1	S	S	S	S	S	S	S		31	35	40	45	50	55	VSP-b	S	S	S	S	S	S	VSP-a	S	S	S	S	S	S	T.phos	S	S	S	S	S	S	Ph.vulg	S	S	S	S	S	S	Ar.VSP	S	S	S	S	S	S	Ar.1A-1	S	S	S	S	S	S	Ar.17A-1	S	S	S	S	S	S		60	65	70	75	80	85	VSP-b	S	S	S	S	S	S	VSP-a	S	S	S	S	S	S	T.phos	S	S	S	S	S	S	Ph.vulg	S	S	S	S	S	S	Ar.VSP	S	S	S	S	S	S	Ar.1A-1	S	S	S	S	S	S	Ar.17A-1	S	S	S	S	S	S		90	95	100	105	110	115	VSP-b	S	S	S	S	S	S	VSP-a	S	S	S	S	S	S	T.phos	S	S	S	S	S	S	Ph.vulg	S	S	S	S	S	S	Ar.VSP	S	S	S	S	S	S	Ar.1A-1	S	S	S	S	S	S	Ar.17A-1	S	S	S	S	S	S
	1	5	10	15	20	25	30																																																																																																																																																																																																																																			
VSP-b	S	S	S	S	S	S	S																																																																																																																																																																																																																																			
VSP-a	S	S	S	S	S	S	S																																																																																																																																																																																																																																			
T.phos	S	S	S	S	S	S	S																																																																																																																																																																																																																																			
Ph.vulg	S	S	S	S	S	S	S																																																																																																																																																																																																																																			
Ar.VSP	S	S	S	S	S	S	S																																																																																																																																																																																																																																			
Ar.1A-1	S	S	S	S	S	S	S																																																																																																																																																																																																																																			
Ar.17A-1	S	S	S	S	S	S	S																																																																																																																																																																																																																																			
	31	35	40	45	50	55																																																																																																																																																																																																																																				
VSP-b	S	S	S	S	S	S																																																																																																																																																																																																																																				
VSP-a	S	S	S	S	S	S																																																																																																																																																																																																																																				
T.phos	S	S	S	S	S	S																																																																																																																																																																																																																																				
Ph.vulg	S	S	S	S	S	S																																																																																																																																																																																																																																				
Ar.VSP	S	S	S	S	S	S																																																																																																																																																																																																																																				
Ar.1A-1	S	S	S	S	S	S																																																																																																																																																																																																																																				
Ar.17A-1	S	S	S	S	S	S																																																																																																																																																																																																																																				
	60	65	70	75	80	85																																																																																																																																																																																																																																				
VSP-b	S	S	S	S	S	S																																																																																																																																																																																																																																				
VSP-a	S	S	S	S	S	S																																																																																																																																																																																																																																				
T.phos	S	S	S	S	S	S																																																																																																																																																																																																																																				
Ph.vulg	S	S	S	S	S	S																																																																																																																																																																																																																																				
Ar.VSP	S	S	S	S	S	S																																																																																																																																																																																																																																				
Ar.1A-1	S	S	S	S	S	S																																																																																																																																																																																																																																				
Ar.17A-1	S	S	S	S	S	S																																																																																																																																																																																																																																				
	90	95	100	105	110	115																																																																																																																																																																																																																																				
VSP-b	S	S	S	S	S	S																																																																																																																																																																																																																																				
VSP-a	S	S	S	S	S	S																																																																																																																																																																																																																																				
T.phos	S	S	S	S	S	S																																																																																																																																																																																																																																				
Ph.vulg	S	S	S	S	S	S																																																																																																																																																																																																																																				
Ar.VSP	S	S	S	S	S	S																																																																																																																																																																																																																																				
Ar.1A-1	S	S	S	S	S	S																																																																																																																																																																																																																																				
Ar.17A-1	S	S	S	S	S	S																																																																																																																																																																																																																																				

TO FIG. 1A.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

METHODS FOR ALTERING THE NUTRITIONAL VALUE OF A PLANT PROTEIN, VEGETATIVE STORAGE PROTEIN, VSP, BY ALTERING THE AMINO ACID CONTENT OF PROTEINS

FIELD OF THE INVENTION

The invention relates to a process for the production of proteins having high nutritional properties. The methods find particular use in the production of plants with increased levels of amino acids having high nutritional properties through the modification of plant genes.

BACKGROUND OF THE INVENTION

Autotrophic organisms can make all of their own amino acids. Other cells utilize many preformed amino acids. Humans and other higher animals require a number of essential amino acids in the diet. These essential amino acids are obtained directly or indirectly by eating plants. These essential amino acids include lysine, tryptophan, threonine, methionine, phenylalanine, leucine, valine and isoleucine.

Constructing proteins with higher nutritional value has been a long-sought goal of scientists. Traditionally, agricultural scientists concentrated on breeding plants with high nutritional yield. Typically, these new varieties were richer in carbohydrates but usually poorer in essential proteins than the wild type varieties from which they were derived.

Seed storage proteins represent up to 90% of total seed protein in seeds of many plants. They are used as a source of nutrition for young seedlings in the period immediately following germination. The genes encoding them are strictly regulated, being expressed in a highly tissue specific and stage specific manner. These genes are almost exclusively expressed in developing seed. Different classes of seed storage proteins may be expressed at different stages in the development of the seed. They are typically stored in membrane bound organelles called protein bodies or protein storage vacuoles.

A related group of proteins, the vegetative storage proteins, have similar amino acid compositions and are also stored in specialized vacuoles. These proteins are generally found in leaves instead of seeds. These proteins are degraded upon flowering, and are thought to serve as a nutritive source for developing seeds.

Cereal grains and legume seeds which are key protein sources for the vegetarian diet are generally deficient in essential amino acids such as methionine, lysine, and threonine. Therefore, there is needed means for improving the nutritional quality of these proteins.

SUMMARY OF THE INVENTION

Compositions and methods for altering the amino acid profiles of proteins without introducing conformational changes into the protein are provided. The method involves preparing a binding partner and/or an interacting molecule which binds to the native protein and using such interacting molecule to select for modified proteins retaining the native conformation.

The method finds particular use in altering the nutritional value of proteins. A plant protein having increased methionine levels is provided. The modified protein retains the conformation of the native protein while having significantly higher levels of methionine.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 shows VSP homologies.

VSP-b (same as VSP β) and VSP-a (same as VSP α): Staswick, P.E., (1988), *Plant Physiol.* **87**, 250-254.

T.phos (tomato acid phosphatase): Erion, J.L., Ballo, B., May, L., Bussell, J., Fox, T.W., & Thomas, S.R., *SwissProt database accession number P27061*.

Ph.vulg (*Phaseolus vulgaris*): Zhon, P-Y., Tanaka, T., Yamauchi, D., & Minamikawa, T. (1997), *Plant Physiol.* **113**, 479-485.

Ar.VSP (*Arabidopsis thaliana*): Yu, D.Y., Quigley, F., & Mache, R.,
EMBL database accession number X79490.

Ar.1A-1, Ar17A-1 (*Arabidopsis thaliana*, floral organs): Utsugi, S.,
Sakamoto, Ogura, Y., Murata, M., & Motoyoshi, F. (1996) *Plant Mol. Biol.* 32,
5 759-765.

Fig. 2 shows proposed VSP β methionine-enriched variants.

Fig. 3A shows the hydropathy index computation for sequence VSP β .

Fig. 3B shows the hydropathy index computation for sequence VSPMet10.

Fig. 3C shows the hydropathy index computation for sequence VSPMet20.

10 Fig. 3D shows the hydropathy index computation for sequence VSPMet30.

Fig. 4 shows the VSP β -met10 sequence.

Fig. 5 shows the colony lift assay to detect protein-protein interactions.

DETAILED DESCRIPTION OF THE INVENTION

15 Proteins having altered amino acid profiles are provided. The proteins can
be designed to be enriched in essential amino acids, including lysine, methionine,
tryptophan, threonine, phenylalanine, leucine, valine and isoleucine relative to
average levels of such amino acids in the native protein. Generally, knowledge
of the three-dimensional (3-D) structure of a given protein allows one to engineer
20 amino acid substitutions in a rational manner so as to effect a desired change in the
property of the protein without compromising the folding process. The present
invention provides methods for increasing the levels of essential amino acids
within a protein while at the same time the altered protein has the conformation of
the native protein.

25 The present invention provides methods for altering the amino acid content
of a protein whose 3-D structure is unknown or unavailable. The method may also
provide an easy method for assessing changes in a protein in which the structure of
the protein is known but tools for confirming conformation of the protein may be
unavailable. The "conformation" of a protein refers to the spatial arrangement of
30 substituent groups of the molecule. The polypeptide chain of a protein has only

one conformation (or a very few) under normal biological conditions of temperature and pH. This, referred to as the "native conformation," confers biological activity. The native conformation is sufficiently stable so that the protein can be isolated and retained in its native state. Therefore, it is important to
5 be able to change the amino acid content of a protein, yet at the same time have the protein retain its biological activity.

The methods of the invention are useful for making amino acid changes within proteins whose conformation is unknown or unavailable. Such proteins include the vegetative storage protein which is believed to play a significant role in
10 supplying amino acids for protein deposition during seed fill, and other proteins of the seed. The methods of the invention may be used to modify the amino acid composition of any protein. Examples of such proteins include but are not limited to wheat endosperm purothionine (Mak and Jones (1976) *Can. J. Biochem.* 22:83J); albumins (Higgins et al. (1986) *J. Biol. Chem.* 261:11124); and
15 methionine rich proteins (Pedersen et al. (1986) *J. Biol. Chem.* 261:6279; Kiriara et al. (1988) *Gene* 71:359; Musumura et al. (1989) *Plant. Mol. Biol.* 12:123).

The methods of the invention comprise altering the amino acid composition of a protein to produce an engineered protein. The engineered protein will retain the conformation and activity of the native protein yet have a modified or altered
20 amino acid content. In this manner, levels of particular amino acids of interest can be increased or decreased. Of particular interest, is to increase the levels or numbers of essential amino acids in the proteins. By essential amino acid is intended, lysine, tryptophan, threonine, methionine, phenylalanine, leucine, valine, isoleucine, and cysteine. However, it is recognized that the amino acid
25 composition can be changed in various ways, as long as the changes do not affect the conformation of the final protein.

The proteins of the invention have been engineered or modified to contain altered amino acid levels. The engineered protein retains the conformation of the native protein. The method involves preparing binding partners and/or interacting
30 molecules to the native protein and utilizing these interacting molecules to

determine whether the engineered protein folds correctly. By "binding partner" or "interacting molecule" is intended a molecule which is capable of binding or interacting with the proteins of interest. Such binding partners or interacting molecules include antibodies, monoclonal antibodies, antibody fragments, proteins, modified proteins, nucleotide sequences, such as aptomers, chemical compounds (e.g. carbohydrates, etc.), or combinations thereof. The interacting molecules also encompass polypeptides that have an intrinsic affinity to the protein of interest, particularly such polypeptides that are capable of binding with the protein of interest to form an oligomeric complex. For example, VSP-alpha binds VSP with high affinity and could be used as an interacting molecule for the altered VSP protein.

Methods for antibody production are known in the art. See, for example, *Antibodies, A Laboratory Manual*, Harlow and Lane (Eds.), Cold spring Harbor Laboratory Press, Coldspring Harbor, NY (1988), and the references cited therein. See also, Radka *et al.* (1983) *J. Immunol.* 128:2804; and Radka *et al.* (1984) *Immunogenetics* 19:63. All of which are herein incorporated by reference.

Once antibodies, preferably monoclonal antibodies, are available which bind to the native protein, such antibodies can be used to select for modified proteins which retain the conformation of the native protein. Strategies to identify residues within a protein that might tolerate amino acid substitution include mutational analysis, secondary structure prediction, homology comparison, and the like. Such strategies can be used to identify amino acids within the protein that will tolerate amino acid substitution.

By mutational analysis is intended mutagenic PCR and DNA shuffling. See, for example, Stemmer, W.P. (1994) *Nature* 370:389-391; and Stemmer, W.P. (1994) *Proc. Natl. Acad. Sci. USA* 91:10747-10751, herein incorporated by reference. Such methods can be used to generate phage display libraries of protein genes containing random mutations. Phage display is an *in vitro* selection technology which allows for a foreign protein or peptide to be displayed on the surface of filamentous phage, linking the phenotype of the phage to its genotype.

Molecular repertoires with sufficient diversity can be generated using such technology. Proteins which exhibit the correct conformation, that is, the native conformation, can be selected for by the ability to bind antibodies recognizing conformational domains of the native protein. See also *Methods in Enzymology*,
5 *Combinatorial Chemistry*, John N. Abelson (Ed.), Vol. 267, Academic Press, Inc., San Diego, CA, herein incorporated by reference. Once correctly-folded protein variants are determined, subsequent isolation and sequencing of the variants reveals the tolerated sites for mutations. Alternatively, correctly folded variants may be identified by other screening or selection methods such as filter lift-assay
10 and ELISA.

Substitutions may also be incorporated at secondary structure prediction sites. Structural features of the protein are important for proper folding. Sequence analysis tools such as the GCG (Wisconsin sequence analysis package, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, WI) and
15 PC/GENE (Oxford Molecular Group, 2105 S. Bascom Avenue, Suite 200, Campbell, CA) can be used to analyze protein sequence for secondary structure features such as helices, sheets and turns. In this manner, it can be determined whether a particular stretch of amino acids may reside on the surface of the protein. Residues on the surface of a protein tolerate substitution more readily than buried
20 residues without compromising the structure of the protein. Utilizing these algorithms, predicted turns and surface regions of the proteins can be made. Therefore, predictions can be made into which regions amino acid substitutions can be made without affecting conformation.

Sites for amino acid substitution can also be determined by homology
25 comparison to other proteins. Nature has tested the tolerance of protein residues to substitution as exemplified in the sequences of proteins such as globins and cytochromes from several different species, members of which have the same fold. See, for example, Hampsey *et al.* (1988) *FEBS Lett.* 231:275; Bashford *et al.* (1987) *J. Mol. Biol.* 196:199; Lesk and Chothia (1980) *J. Mol. Biol.* 136:225.

In designing proteins of the invention, hydrophobic residues, such as alanine, cysteine, valine, isoleucine, leucine, methionine, phenylalanine, and tryptophan may be substituted for one another without undue perturbation of the structure. Such residues generally occur in the hydrophobic core of the protein.

5 See, Bowie *et al.* (1990) *Science* 247:1306-1310; and Baldwin and Matthews (1994) *Curr. Opin. Biotech* 5:396-402. See also Ladunga and Smith (1997) *Prot. Eng.* 10:187-196, herein incorporated by reference. Generally, residues that substitute for one another in related sequences do so by conserving the physico-chemical properties of the residue and folding of the protein thus conserving the 3-D structure of the protein.

10

Therefore, the protein to be modified can be compared with homologous proteins. Amino acids that are critical to the function and/or folding of the protein would be expected to be conserved over time. Therefore, predictions can be made as to which amino acids can be substituted without affecting the conformation or folding of the protein.

15

Such selected amino acid substitutions can be made by DNA sequencing, site-directed mutagenesis, or other methods which substitute one amino acid with any other amino acid.

Once the amino acid substitutions have been made and the conformation confirmed by antibody binding, the protein can be expressed using known expression systems. Where necessary, the DNA encoding the protein can be synthesized using known techniques. Likewise, the nucleotide sequence encoding the protein can be contained within expression cassettes.

20

Utilizing the methods of the invention, proteins can be constructed which have increased nutritional quality. That is, the essential amino acid content within the protein can be increased to represent at least about 5-about 10%, preferably at least about 10-about 20%, more preferably at least about 20-about 40% of the total amino acid content in the protein.

25

In the same manner, the amino acid content of a subject protein can be altered to include at least about 10% amino acid substitutions, additions or

30

deletions, about 20% or even up to about 30% to about 40%. It is recognized that the limitation will be the activity of the altered. The present invention provides a convenient and ready mechanism to test the activity of the protein by its ability to bind the interacting molecule.

5 For convenience for expression in plants, the nucleic acid encoding the modified peptides or proteins of the invention can be contained within expression cassettes. The expression cassette will comprise a transcriptional initiation region linked to the nucleic acid encoding the peptide of interest. Such an expression cassette is provided with a plurality of restriction sites for insertion of
10 the gene or genes of interest to be under the transcriptional regulation of the regulatory regions.

The transcriptional initiation region, the promoter, may be native or homologous or foreign or heterologous to the host, or could be the natural sequence or a synthetic sequence. By foreign is intended that the transcriptional
15 initiation region is not found in the wild-type host into which the transcriptional initiation region is introduced.

The transcriptional cassette will include the in 5N-3N direction of transcription, a transcriptional and translational initiation region, a DNA sequence of interest, and a transcriptional and translational termination region
20 functional in plants. The termination region may be native with the transcriptional initiation region, may be native with the DNA sequence of interest, or may be derived from another source. Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. See also, Guerinneau *et al.*,
25 (1991) *Mol. Gen. Genet.* 262:141-144; Proudfoot (1991) *Cell* 64:671-674; Sanfacon *et al.* (1991) *Genes Dev.* 5:141-149; Mogen *et al.* (1990) *Plant Cell* 2:1261-1272; Munroe *et al.* (1990) *Gene* 91:151-158; Ballas *et al.* 1989) *Nucleic Acids Res.* 17:7891-7903; Joshi *et al.* (1987) *Nucleic Acid Res.* 15:9627-9639.

Where appropriate, the gene(s) expressing the modified proteins may be
30 optimized for increased expression in the transformed plant. In this manner, the

sequences can be synthesized using monocot, dicot or particular plant; i.e. maize, soybean, sorghum, wheat, etc., preferred codons for improved expression.

Methods are available in the art for synthesizing plant preferred genes. See, for example, U.S. Patent Nos. 5,380,831, 5,436, 391, and Murray et al. (1989)

5 *Nucleic Acids Res.* 17:477-498, herein incorporated by reference.

The expression cassettes may additionally contain 5' leader sequences in the expression cassette construct. Such leader sequences can act to enhance translation. Translation leaders are known in the art and include: picornavirus leaders, for example, EMCV leader (Encephalomyocarditis 5' noncoding region) (Elroy-Stein, O., Fuerst, T.R., and Moss, B. (1989) *PNAS USA*, 86:6126-6130);
10 potyvirus leaders, for example, TEV leader (Tobacco Etch Virus) (Allison *et al.* (1986); MDMV leader (Maize Dwarf Mosaic Virus); *Virology*, 154:9-20), and human immunoglobulin heavy-chain binding protein (BiP), (Macejak, D.G., and P. Sarnow (1991) *Nature*, 353:90-94; untranslated leader from the coat protein
15 mRNA of alfalfa mosaic virus (AMV RNA 4), (Jobling, S.A., and Gehrke, L., (1987) *Nature*, 325:622-625; tobacco mosaic virus leader (TMV), (Gallie, D.R. *et al.* (1989) *Molecular Biology of RNA*, pages 237-256; and maize chlorotic mottle virus leader (MCMV) (Lommel, S.A. *et al.* (1991) *Virology*, 81:382-385). See also, Della-Cioppa *et al.* (1987) *Plant Physiology*, 84:965-968.

20 Other methods known to enhance translation can also be utilized, for example, introns, and the like.

The expression cassettes may contain one or more than one nucleic acid sequences to be transferred and expressed in the transformed plant. Thus, each nucleic acid sequence will be operably linked to 5' and 3' regulatory sequences.

25 Alternatively, multiple expression cassettes may be provided.

Generally, the expression cassette will comprise a selectable marker gene for the selection of transformed cells. Selectable marker genes are utilized for the selection of transformed cells or tissues. Such selectable marker genes are known in the art. See generally, G. T. Yarranton (1992) *Curr. Opin. Biotech.*,
30 3:506-511; Christopherson *et al.* (1992) *Proc. Natl. Acad. Sci. USA*, 89:6314-

6318; Yao *et al.* (1992) *Cell*, 71:63-72; W. S. Reznikoff (1992) *Mol. Microbiol.*, 6:2419-2422; Barkley *et al.* (1980) *The Operon*, pp. 177-220; Hu *et al.* (1987) *Cell*, 48:555-566; Brown *et al.* (1987) *Cell*, 49:603-612; Figge *et al.* (1988) *Cell*, 52:713-722; Deuschle *et al.* (1989) *Proc. Natl. Acad. Sci. USA*, 86:5400-5404; Fuerst *et al.* (1989) *Proc. Natl. Acad. Sci. USA*, 86:2549-2553; Deuschle *et al.* (1990) *Science*, 248:480-483; M. Gossen (1993) Ph.D. Thesis, University of Heidelberg; Reines *et al.* (1993) *Proc. Natl. Acad. Sci. USA*, 90:1917-1921; Labow *et al.* (1990) *Mol. Cell Bio.*, 10:3343-3356; Zambretti *et al.* (1992) *Proc. Natl. Acad. Sci. USA*, 89:3952-3956; Baim *et al.* (1991) *Proc. Natl. Acad. Sci. USA*, 88:5072-5076; Wyborski *et al.* (1991) *Nuc. Acids Res.*, 19:4647-4653; A. Hillenand-Wissman (1989) *Topics in Mol. and Struc. Biol.*, 10:143-162; Degenkolb *et al.* (1991) *Antimicrob. Agents Chemother.*, 35:1591-1595; Kleinschmidt *et al.* (1988) *Biochemistry*, 27:1094-1104; Gatz *et al.* (1992) *Plant J.*, 2:397-404; A. L. Bonin (1993) PhD Thesis, University of Heidelberg; Gossen *et al.* (1992) *Proc. Natl. Acad. Sci. USA*, 89:5547-5551; Oliva *et al.* (1992) *Antimicrob. Agents Chemother.*, 36:913-919; Hlavka *et al.* (1985) *Handbook of Exp. Pharmacology*, 78; Gill *et al.* (1988) *Nature* 334:721-724; DeBlock *et al.* (1987) *EMBO J.*, 6:2513-2518; DeBlock *et al.* (1989) *Plant Physiol.*, 91:691-704; Fromm *et al.* (1990) 8:833-839; Gordon-Kamm *et al.* (1990) 2:603-618. Such disclosures are herein incorporated by reference.

The nucleotide sequences of interest of this invention can be introduced into the genome of the desired host organism in a variety of techniques known in the art. For the purposes of this invention, it will be appreciated to those skilled in the art that any conventional transformation vector may be used as long as it is capable of transforming the organism of choice and it does not have restriction sites in common with those comprising the final master insertion cassette. Hence, the detailed experimental description of transformation vectors is given by way of illustration only.

Vector systems are known for the transformation of yeast and bacterial cells. For yeast, these include but are not limited to autonomously replicating

plasmids (see, for example, Stearns *et al.* (1990) *Methods Enzymol.* 185:280-297); 2-micron circle yeast DNA sequences (see, for example, Hollenberg (1982) *Curr. Topics Microbiol. Immunol.* 96:119-144; Broach (1983) *Methods Enzymol.* 101:307-325; MacKay (1983) *Methods Enzymol.* 101:325-343; Armstrong (1989) *BioTechnology* 13:165-192; Rose (1990) *Methods Enzymol.* 185:234-279); linearized vector DNA (see, for example, see, for example, Takita *et al.* (1997) *Yeast* 13:763-768); artificial chromosome vectors (Burke (1987) *Science* 236:806-812); restriction site bank plasmids (Davison (1987), U.S. Patent No. 4,657,858, and *Methods Enzymol.* 153: 34-54); delta-integration vectors (see, for example, Lee and Da Silva (1997) *Biotechnol. Prog.* 13:368-373); and *Agrobacterium*-based vectors (see, for example, Bundock *et al.* (1995) *EMBO J.* 14:3206-3214; Piers *et al.* (1996) *Proc. Natl. Acad. Sci. USA* 93:1613-1618; Risseuw *et al.* (1996) *Mol. Cell. Biol.* 16:5924-5932); and Shuttle Vectors (see, for example, Schneider (1991) *Methods Enzymol* 194:373-388; Singh (1997) *Methods Mol. Biol.* 62:113-130). See generally Hinnen (1980) *Curr. Topics Microbiol. Immunol.* 96:101-117; Nombela (1985) *Revis. Biol. Cel.* 4:1-25; Parent (1985) *Yeast* 1(2):83-138; West (1988) *BioTechnology* 10:387-404; Schena (1991) *Methods Enzymol.* 194:389-398; Schneider (1991) *Methods Enzymol.* 194:373-388; and Singh (1997) *Methods. Mol. Biol.* 62:113-130.

Vector systems used for bacterial transformation include, but are not limited to, yeast shuttle vectors (see, for example, Ward (1990) *Nucleic Acids Res.* 18(17):5319; Strathern (1991) *Methods Enzymol.* 194:319-329; Soni (1992) *Nucleic Acids Res.* 20(21) 5852; Nacken (1994) *Nucleic Acids Re.* 22:1509-1510; Wehmeier (1995) *Gene* 165:149-150); pBR322 and related plasmids such as pBR327 and pKC7 (see, for example, Rao and Rogers (1979) *Gene* 7:79-82; Talmadge and Gilbert (1980) *Gene* 12:235-241; Smith *et al.* (1995) *Microbiology* 141(pt. 1):181-188); pATH vectors (see, for example, Koerner *et al.* (1991) *Methods in Enzymol.* 194:477-490); yeast plasmids (see, for example, Marcil (1992) *Nucleic Acids Res.* 20:917); and natural replicon ColEI and related plasmids such as P15A, F, RSF1010, and R616 (see, for example, Muhlenhoff

and Chauvat (1996) *Mol. Gen. Genet.* 252:93-100; Sakai and Komano (1996) *Biosci. Biotechnol. Biochem.* 60:377-382; Lee and Henk (1997) *Vet. Microbiol.* 54:369-374); herein incorporated by reference.

5 A number of vector systems are also known for the introduction of foreign or native genes into mammalian cells. These include SV40 virus (see, for example, Okayama *et al.* (1985) *Molec. Cell. Biol.* 5:1136-1142); Bovine papilloma virus (see, for example, DiMaio *et al.* (1982) *Proc. Natl. Acad. Sci. USA* 79:4030-4034); adenovirus (see, for example, Morin *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:4626; Yifan *et al.* (1995) *Proc. Natl. Acad. Sci. USA* 92:1401-1405; Yang *et al.* (1996) *Gene Ther.* 3:137-144; Tripathy *et al.* (1996) *Nat. Med.* 2:545-550; Quantin *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:2581-2584; Rosenfeld *et al.* (1991) *Science* 252:431-434; Wagner (1992) *Proc. Natl. Acad. Sci. USA* 89:6099-6103; Curiel *et al.* (1992) *Human Gene Therapy* 3:147-154; Curiel (1991) *Proc. Natl. Acad. Sci. USA* 88:8850-8854; LeGal LaSalle *et al.* (1993) *Science* 259:590-599); Kass-Eisler *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:11498-11502); adeno-associated virus (see, for example, Muzyczka *et al.* (1994) *J. Clin. Invest.* 94:1351; Xiao *et al.* (1996) *J. Virol.* 70:8098-8108); herpes simplex virus (see, for example, Geller *et al.* (1988) *Science* 241:1667; Huard *et al.* (1995) *Gene Therapy* 2:385-392; U.S. Patent No. 5,501,979);
10 retrovirus-based vectors (see, for example, Curran *et al.* (1982) *J. Virol.*, 44:674-682; Gazit *et al.* (1986) *J. Virol.*, 60:19-28; Miller (1992) *Curr. Top. Microbiol. Immunol.* 158:1-24; Cavanaugh *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:7071-7075; Smith *et al.* (1990) *Mol. Cell. Biol.* 10:3268-3271); herein incorporated by reference.

15 25 Methods of the present invention can be used to facilitate assembly of nucleotide sequences of interest for transformation of any plant. In this manner, genetically modified plants, plant cells, plant tissue, seed, and the like can be obtained. The transformation vector and hence method of transformation chosen will depend on the type of plant or plant cell, i.e. monocot or dicot, targeted for
30 transformation. Suitable methods of transforming plant cells include

microinjection (Crossway *et al.* (1986) *Biotechniques* 4:320-334); electroporation (Riggs *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 83:5602-5606); *Agrobacterium*-mediated transformation (Hinchey *et al.* (1988) *Biotechnology* 6:915-921); direct gene transfer (Paszkowski *et al.* (1984) *EMBO J.* 3:2717-2722); and ballistic particle acceleration (see, for example, Sanford *et al.* U.S. Patent 4,945,050; WO91/10725 and McCabe *et al.* (1988) *Biotechnology* 6:923-926). Also see, Weissinger *et al.* (1988) *Annual Rev. Genet.* 22:421-477; Sanford *et al.* (1987) *Particulate Science and Technology* 5:27-37 (onion); Christou *et al.* (1988) *Plant Physiol.* 87:671-674 (soybean); McCabe *et al.* (1988) *BioTechnology* 6:923-926 (soybean); Datta *et al.* (1990) *Biotechnology* 8:736-740 (rice); Klein *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:4305-4309 (maize); Klein *et al.* (1988) *BioTechnology* 6:559-563 (maize); WO91/10725 (maize); Klein *et al.* (1988) *Plant Physiol.* 91:440-444 (maize); Fromm *et al.* (1990) *BioTechnology* 8:833-839; and Gordon-Kamm *et al.* (1990) *Plant Cell* 2:603-618 (maize); Hooydaas-Van Slogteren and Hooykaas (1984) *Nature (London)* 311:763-764; Bytebier *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:5345-5349 (Liliaceae); De Wet *et al.* (1985) In *The Experimental Manipulation of Ovule Tissues*, ed. G.P. Chapman *et al.*, pp. 197-209 (Longman, N.Y.) (pollen); Kaeppler *et al.* (1990) *Plant Cell Reports* 9:415-418; and Kaeppler *et al.* (1992) *Theor. Appl. Genet.* 84:560-566 (whisker-mediated transformation); D' Halluin *et al.* (1992) *Plant Cell* 4:1495-1505 (electroporation); Li *et al.* (1993) *Plant Cell Reports* 12:250-255, and Christou and Ford (1995) *Annals of Botany* 75:407-413 (rice); Osjoda *et al.* (1996) *BioTechnology* 14:745-750 (maize via *Agrobacterium tumefaciens*); all of which are herein incorporated by reference.

The following examples are offered by way of illustration and not by way of limitation.

EXPERIMENTAL

Three complementary strategies, namely, mutational analysis, secondary structure prediction, and homology comparison (see below) have been used to

identify amino acids within VSP β (vegetative storage protein) that might tolerate methionine substitution. Together, results from these strategies facilitated the design of three VSP variants with increasing methionine content.

5 1. ***Mutational analysis***

The simple premise behind this strategy was that if one prepared monoclonal antibodies that recognized the wild-type VSP, then these same antibodies would, if the mutant proteins folded correctly, also recognize the engineered proteins. As a first step, therefore, mice were injected with VSP purified from soybean leaves, and a panel of 21 monoclonal antibodies recognizing wild-type VSP has been characterized by ELISA. These antibodies also recognize VSP α expressed and purified from *Pichia pastoris*.

The following two approaches can be implemented to generate either random or "semi-rational" mutations in VSP β . Mutagenic PCR and DNA shuffling (Stemmer, W.P. (1994) *Nature* 370, 389-391; Stemmer, W.P. (1994) *Proc. Natl. Acad. Sci. USA* 91, 10747-10751) can be used to generate phage display libraries of VSP β genes containing random mutations. Since these mutations could alter the structure of VSP, correctly-folded variants can be selected for by their ability to bind a set of monoclonal antibodies recognizing different conformational domains of wild-type VSP. Likewise, correctly-folded variants can be selected by their abilities to homo-heterodimerize. Correctly-folded VSP variants (*i.e.*, those retaining the ability to bind VSP-specific conformational antibodies and homo/heterodimerize) can be selected by phage display technology or screened using a filter lift assay (see methods). Subsequent isolation and sequencing of these variants reveals the tolerated mutations. Amino acid substitutions which do not compromise the VSP β structure may be good candidates for site-directed methionine substitutions.

In addition to this "random" approach, a method for the "semi-rational" incorporation of methionines into VSP was developed. Although the 3-D structure of VSP is uncertain, secondary structure prediction of the protein (see strategy 2

below) allowed "semi-rational" methionine substitutions. Analysis of VSP β homology with tomato acid phosphatase, a protein with 45% identity to VSP β , as well as other homologs allowed additional methionine substitutions (see strategy 3 below). Two methods were designed by which to introduce these substitutions.

5 The first method involves DNA shuffling in the presence of excess methionine-encoding oligos which, by protein secondary structure predictions, are complementary to multiple regions of the VSP β gene corresponding to protein loops. The second novel method employed overlap PCR of segments of the VSP β gene corresponding to protein loops which have been amplified with the
10 methionine-encoding oligos. The methods by which these oligos (corresponding to, for example, twenty-two different methionine substitutions) are introduced into VSP β result in the production of a library of phage-displayed VSP variants; theoretically each variant contains zero to twenty-two additional methionines. Subsequent phage display and biopanning of these libraries against VSP-specific
15 monoclonal antibodies can lead to the identification of residues in VSP which can accommodate methionine without significantly altering the structure of the protein.

A VSP β mutant library was made by error prone PCR methodology (see below). From this pool of mutants, a filter lift assay (see methods) was performed to identify properly-folded mutant VSP β based on the ability to bind to either
20 VSP α or a VSP-specific monoclonal antibody. Using VSP α as the antigen in a filter lift assay (Fig. 5) 18 out of 50 VSP β variants tested bound VSP α . Sequence analysis of 15 of these variants revealed a total of 84 point mutations which correlate with 58 AA substitutions and 25 silent mutations. Together these represent 51 different residues within the 218 AA VSP β .

25

2. *Secondary structure prediction*

Structural features of a protein are very important for proper folding. Sequence analysis tools such as the GCG (Wisconsin Sequence Analysis Package, Genetic Computer Group, University Research Park, 575 Science Drive, Madison,
30 WI) and PC/GENE (Oxford Molecular Group, 2105 S. Bascom Avenue, Suite 200,

Campbell, CA) were used to analyze the VSP β sequence for secondary structure features such as helices, sheets and turns and for determining whether a particular stretch of amino acids might reside on the surface of the protein. Residues on the surface of a protein would likely tolerate substitution more readily than a buried residue without compromising the structure of the protein. Using these algorithms, numerous predicted turns and surface regions of the protein were identified. Many of these regions are expected to tolerate methionine substitution. For example residues at positions 25, 30, 32, 37, 44, 65, 67, 102, 121, 130, 160, 163, 164, 169, 198, 202, and 207 in VSP β occur in predicted turn regions and were substituted with Met (Table 1).

3. *Homology comparison*

Over time, nature has tested the tolerance of protein residues to substitution, and this is exemplified in the sequences of proteins such as globins and cytochromes from several different species, members of which have the same fold (Hampsey, M.D., Das, G., Sherman, F. (1988) *FEBS Lett.* 231,275; Bashford, D., Chothia, C. & Lesk, A.M. (1987) *J. Mol. Biol.* 196, 199; Lesk, A.M. & Chothia, C. (1980) *J. Mol. Biol.* 136,225). These and other studies have demonstrated that hydrophobic residues (such as Ala, Val, Ile, Leu, Met, Phe and Trp) almost always occur in the hydrophobic core of the protein and that they may substitute for each other without undue perturbation of the structure (Bowie, J.U., Reidhaar-Olson, J.F., Lim, W.A., & Sauer, R.T. (1990) *Science* 247, 1306-1310; Baldwin, E.P., & Matthews, B.W. (1994) *Curr. Opin. Biotech.* 5, 396-402). Indeed, it has been observed that "Residue positions that can accept a number of different side chains, including charged and highly polar residues, are almost certain to be on the protein surface. Bowie *et al.* (1990) *Science* 247:1306-1310, have Residue positions that remain hydrophobic, whether variable or not, are likely to be buried within the structure". Furthermore, in a recent comprehensive analysis of substitution patterns in several databases of multiply aligned protein sequences, Ladunga and Smith (1997) *Prot. Eng.* 10:187-196, have concluded that the overall

emphasis is on the preservation of three dimensional structure of the protein and that residues that substitute for each other in related sequences do so by conserving the physico-chemical properties of the residue and the folding of the protein. In the case of VSP, this evolutionary data was utilized by comparing the homology of VSP β with six homologous proteins (Fig. 1). Amino acids that are critical to the function and/or folding of a protein would be expected to be conserved over time. For example, cysteine 7 and 29 are conserved in all seven of the homologous proteins aligned in Fig. 1. These residues are involved in forming a disulfide bond that may be expected to be of importance to the structure of the protein.

In summary, analysis of the VSP β sequence with its homologs led to the identification of 31 residues (out of 218 amino acids) that in all likelihood will tolerate methionine substitution.

Engineering VSP β for increased methionine

Rational

Wild-type VSP β contains 1.4% methionine. Using the three strategies described, three different VSP β variants with increasing amounts of methionine have been proposed (9.6%, 14.2%, 17.9%, Fig. 2). The overall amino acid composition in each of these constructs is presented in Table 2. Construct VSP β -met20 (14.2% Met) contains the same 18 Met substitutions as the VSP β -met10 derivative plus an additional 11 Met residues. Likewise VSP β -met30 contains the same 29 Met substitutions as VSP β -met20 plus an additional 7 Met residues. Mutational analysis of VSP β resulted in the mutation of 51 different amino acids out of the 218 amino acid protein. Although these mutations were not methionine substitutions, the types of tolerated substitutions were examined for their relevance to substitution to a hydrophobic amino acid. For example, positions 50, 67, 93, 127, 150, and 164 tolerated mutation to a hydrophobic amino acid (Table 1). Therefore, it is possible that this same position might tolerate substitution to methionine. Positions 62, 67, 76, 127, and 164 are hydrophobic amino acids in

VSP β - wild type. The observation that these positions tolerate substitution at all suggests they would more readily tolerate a conservative substitution (*i.e.*, hydrophobic amino acid to hydrophobic amino acid, Table 1). Since residues 32, 50, 65, 67, 76, 93, 127, 150, 160, and 202 allowed non-conservative mutations, it is possible that these positions would tolerate mutation to methionine (Table 1). In every case where these amino acids were not changed from or to a hydrophobic amino acid in the mutational analysis, at least one additional strategy (*i.e.*, secondary structure or homology comparison) was used to rationalize methionine substitution at the particular position. In summary, in the three methionine enriched constructs proposed, 12 residues (out of a total of 36) were selected based at least in part on mutational analysis. More specifically, mutational analysis indicated 6/18 methionine substitutions in construct VSP β -met10, 9/29 in construct VSP β -met20, and 12/36 in VSP β -met30 (Table 1). As mentioned, mutational analysis revealed 51 different positions within VSP β tolerant to substitutions. Interestingly, 25/51 (49%) of the mutated positions are located in regions of the protein predicted to exist as turns, 17/51 (33%) in helices, and 9/51 (18%) in β -sheets. These percentages are significantly different from the predicted distribution of turns (25%), helices (25%) and β -sheets (50%), indicating that, as expected, the regions of the protein most likely to be located on the surface (*e.g.*, turns) can more readily accommodate substitutions without compromising the structure of the protein. This suggests the importance of protein secondary structure prediction as one of the strategies utilized in the identification of residues for methionine substitution.

Since protein turns are generally more surface-exposed regions that do not contribute greatly to the overall structure of the protein, these regions were targeted for methionine substitution. In fact, out of the 36 positions selected for methionine substitution, 17 (47.2%) are predicted to occur in turns. In contrast, because β -sheets are protein structural elements that generally occur at the core of the protein, these regions were avoided in selecting sites for methionine substitution. Out of the 36 positions selected for methionine substitution, only 7 (19.4%) are predicted

to occur in β -sheets. Nearly all of these residues were hydrophobic in wild-type VSP β and were thought to tolerate methionine based upon the homology comparison strategy. Additionally, 12 (33.3%) of the residues selected for methionine substitution in the three constructs are predicted to occur in helices. In
5 summary, secondary structure prediction is the strategy responsible, at least in part, for 17/36 sites targeted for methionine substitution. More specifically, secondary structure prediction correlates with the selection of 7/18, 14/29, and 17/36 amino acids for methionine substitution in constructs VSP β -met10, VSP β -met20, and VSP β -met30, respectively (Table 1).

10 Homology comparison was a very informative strategy in selecting residues that might tolerate methionine substitution. Accordingly, methionine substitutions in VSP β were made by adhering to the following rules and also summarized in Table 1:

(a) Conserved residues (highlighted in blue in Fig. 1) were defined as
15 those residues occurring in more than 5 of the 7 homologs. These were not targeted for substitution. The exceptions were: at residue numbers 19, 37, 146 and 179 (one of the homologs contained a methionine residue); at positions 67, 80, 130 and 169 (conserved hydrophobic amino acid exchanges observed in at least one sequence) and at position 50 (non-conservative changes from Asn to Ser/Cys in
20 two sequences).

(b) Similarly, non-conserved positions were defined as those containing residues with different side-chain properties. Several positions in VSP β were correlated with non-conservative amino acids in the homologs (*e.g.*, 5, 19, 25, 30, 37, 44, 60, 62, 65, 67, 72, 76, 80, 90, 97, 102, 121, 127, 130, 135, 142, 146, 150,
25 164, 169, 179, 189, 198, 202, 207, and 217). Such residues likely reside on the surface/turns of the protein and were considered less important for protein function and/or folding and therefore targeted for substitution with methionine.

(c) In addition, some positions in which at least one other hydrophobic amino acid was observed among homologs (*e.g.*, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 12,
30 13, 14, 15, 16, 17, 18, 19, 20, 21, 25, 30, 37, 44, 60, 62, 65, 67, 72, 76, 90, and 97)

were also expected to tolerate substitution to the hydrophobic amino acid methionine. Exceptions to this were cases in which the hydrophobic amino acid was completely conserved in all 6 homologs (*e.g.*, Val 49, Leu 77, Leu 110, Leu 114, Leu 145, Ile 157, Leu 158, Ile 186, Val 187, Leu 197 and Leu 210). In these cases, the possibility that the specific hydrophobic amino acid in the wild-type protein may be playing a role critical for the proper structure and/or function of the protein was considered. To avoid disturbing this possible role, the substitution of any residue that is completely conserved in all 6 homologs examined was not proposed.

- 10 (d) Six residues within VSP β that were expected to tolerate methionine substitution were identified based on the presence of methionine in analogous positions in homologs (*e.g.*, 19, 37, 44, 146, 179, and 202).

A few additional considerations were observed in selecting amino acids that might tolerate methionine substitution.

- 15 (e) We avoided altering histidine residues due to their potential importance in phosphatase activity of VSP β (Table 2 and DeWald, D.B., Mason, H.S., & Mullet, J.E. (1992) *J. Biol. Chem.* 267, 15958-15964).

- (f) Since VSP β is a glycoprotein, this feature may be important for the stability and/or function of the protein, substitution of potential glycosylation sites was avoided (*e.g.*, Asn 94).

- 20 (g) In addition, wherever possible, charged residues such as Lys, Arg, Glx, Asx were left untouched to preserve the hydrophobic/hydrophilic balance of the protein (Table 2 and Fig. 3A-D). While wild-type VSP β has a calculated charge of -4, VSP β -met10, VSP β -met20, and VSP β -met30 have calculated charges of -7, -7, and -5, respectively.

25 As a strategy, homology comparison facilitated, at least in part, the selection of 31/36 of the residues proposed for methionine substitution. These selections correlate with 18/18, 28/29, and 31/36 residues for constructs VSP β -met10, VSP β -met20, and VSP β -met30, respectively (Table 1).

Several of the amino acids selected for methionine substitution in the three constructs resulted from more than one strategy. In fact, the majority (20/36) of the targeted residues resulted from at least two strategies, with a few (4/36) resulting from all three strategies.

5

Experimental results

A synthetic gene for methionine enriched VSP β -met10 has been constructed. This synthetic gene differs from wild-type VSP β in that it encodes eighteen additional methionines (Fig. 4). Also, a few silent point mutations were introduced into this construct to create unique restriction sites. To test whether the proposed VSP β -met10 gene was correctly folded, the construct was cloned into the phagemid vector pCANTAB-5E and the abilities of the expressed proteins to bind VSP-specific conformational monoclonal antibodies in a filter lift assay were compared. The results indicate that the VSP β -met10 gene was able to bind the same antibodies as wild-type VSP β . This suggests that VSP β -met10 may be correctly folded in an *E. coli* secretion system.

10

15

20

Together, these interdisciplinary approaches should not only result in the engineering of a nutritionally-enhanced VSP, but also provide clues to the structure of VSP - a protein for which no 3D structure is available. This approach is applicable to any protein of interest.

Methods

1. Random mutation of vegetative storage protein (VSP β) by error-prone PCR

25

The VSP β gene was amplified by mutagenic PCR using primers flanking the gene.

<u>Reaction 1</u>	<u>Reaction 2</u>	<u>Reaction 3</u>	<u>Reaction 4</u>
10mM Tris-HCl	10mM Tris-HCl	10mM Tris-HCl	10mM Tris-HCl
50mM KCl	50mM KCl	50mM KCl	50mM KCl
9.5mM MgCl ₂	9.mM mgCl ₂	9.mM mgCl ₂	9.mM mgCl ₂
0.5mM MnCl ₂	0.5mM MnCl ₂	0.5mM MnCl ₂	0.5mM MnCl ₂
5 µg/ml BSA	5 µg/ml BSA	5 µg/ml BSA	5 µg/ml BSA
600pmol VSP template	600pmol VSP template	600pmol VSP template	600pmol VSP template
0.1 µm each primer	0.1 µm each primer	0.1 µm each primer	0.1 µm each primer
2mM dATP	200µM dATP	200µM dATP	200µM dATP
200µM dCTP	2mM dCTP	200µM dCTP	200µM dCTP
200µM dGTP	200µM dGTP	2mM dGTP	200µM dGTP
200µM dTTP	200µM dTTP	200µM dTTP	2mM dTTP
2 Units Taq Pol	2 Units Taq Pol	2 Units Taq Pol	2 Units Taq Pol
1 cycle (1 min. at 95EC, 1 min. at 51EC, 3 min. at 72EC)			
16 cycles (1 min. at 91EC, 1 min. at 51EC, 3 min. at 72EC)			
1 cycle (1 min. a 91EC, 1 min. at 51EC, 5 min. at 72EC)			

The products of these four reactions were pooled, and the band corresponding to the mutagenized VSPβ gene was purified from an agarose gel, digested with *Sfi*I and *Not*I and cloned into the phagemid vector pCANTAB-5E.

2. Filter lift assay

Fifty *E. coli* colonies containing randomly mutated VSPβ genes were picked as small patches to an SB agar plate containing glucose and ampicillin.

Patches were allowed to grow overnight at 37EC and were then transferred to a nitrocellulose filter. On the surface of an SB agar plate containing ampicillin and IPTG, this filter was placed on top (cell-side up) of a separate blocked filter to which the antigen (*e.g.*, VSP α) had been coated. During an overnight incubation at 30EC, the cells expressed the VSP β variant they encoded. These proteins were able to diffuse through the top filter and, if correctly folded, bind the antigen-coated filter below. The next day, the antigen-coated filter was washed with PBS-0.05% tween and incubated with HRP/anti-e tag conjugate. Since the VSP β mutants are cloned into the pCANTAB-5E vector which fuses a C-terminal epitope tag (e-tag) to the VSP β protein variants, bound proteins were detected by this antibody in combination with enhanced chemiluminescence detection.

Table 1

PROPOSED METHIONINE SUBSTITUTION

		Homology comparison				
VSPB position		Mutational Analysis ¹	Secondary Structure ²	Original A.A. hydrophobic?	Met in homolog? ³	# homologs hydrophobic ⁴
<i>Construct 1 (9.6% Met)</i>						
5	V	-	-	Y	-	3 of 6
19	I	-	-	Y	Y-Ar. VSP	6 of 6
30	V	-	T	Y	-	2 of 6
37	I	-	T	Y	Y-T, phos	6 of 6
44	I	-	T	Y	Y-T, phos	1 of 6
60	R	-	-	-	-	5 of 6
62	V	V-A	-	Y	-	6 of 6
67	I	I-T,L	T	Y	-	5 of 6
72	I	-	-	Y	-	6 of 6
76	V	V-G	-	Y	-	5 of 6
121	L	-	T	Y	-	6 of 6
127	I	I-T,L	-	Y	-	3 of 6
146	K	-	-	-	Y-T, phos	1 of 6
164	I	I-V	T	Y	-	3 of 5
179	L	-	-	Y	Y-T, phos	6 of 6
189	I	-	-	Y	-	2 of 6
202	R	R-G,T	T	-	Y-T, phos	1 of 6
217	I	-	-	Y	-	5 of 6
<i>Construct 2-additional substitution (14.7% Met)</i>						
32	P	P-Q	T	-	-	0 of 6
65	N	N-S	T	-	-	3 of 6
90	V	-	-	Y	-	2 of 6
97	L	-	-	Y	-	1 of 6
102	V	-	T	Y	-	5 of 6
130	L	-	T	Y	-	6 of 6
135	L	-	-	Y	-	1 of 5
150	F	F-S,I,L	-	-	-	3 of 6
169	L	-	T	Y	-	5 of 6
198	L	-	T	Y	-	5 of 6
207	T	-	T	-	-	3 of 6
<i>Construct 3-additional substitutions (17.9% Met)</i>						
25	I	-	T	Y	-	6 of 6
50	N	N-I	-	-	-	0 of 6
80	I	-	-	Y	-	6 of 6
93	F	F-V	-	-	-	0 of 6
142	E	-	-	-	-	3 of 6
160	D	D-Y	T	-	-	0 of 6
163	L	-	T	Y	-	0 of 6

¹ Amino acid substitution observed in the mutational analysis. For example, at position 62,a valine to alanine substitution was observed

² "T" indicates turn predicted by secondary structure analysis of VSPB.

³ "Y" indicates the presence of Methionine in the designated VSP homolog.

⁴ Includes only aliphatic hydrophobic amino acids such as Leu, Ile, Val, and Met.

Table 2

Amino Acid Composition of VSPB-WT and Methionine-Enriched Variants

	<u>VSPB-WT</u>	<u>VSPB-Met10</u>	<u>VSPB-Met20</u>	<u>VSPB-Met30</u>
Ala	13	13	13	13
Arg	11	9	9	9
Asn	14	14	13	12
Asp	11	11	11	10
Cys	2	2	2	2
Glu	6	6	6	6
Glu	19	19	19	18
Gly	13	13	13	13
His	7	7	7	7
Ile	14	6	6	4
Leu	20	18	13	12
Lys	15	14	14	14
MET	3 (1.4%)	21 (9.6%)	32 (14.7%)	39 (17.9%)
Phe	12	12	11	10
Pro	9	9	8	8
Ser	13	12	12	12
Thr	10	10	9	9
Trp	3	3	3	3
Tyr	12	12	12	12
Val	11	7	5	5
Total	218	218	218	218

All publications and patent applications mentioned in the specification are indicative of the level of those skilled in the art to which this invention pertains. All publications and patent applications are herein incorporated by reference to the

same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be
5 obvious that certain changes and modifications may be practiced within the scope of the appended claims.

CLAIMS:

1. A method for altering amino acid composition of a native protein of interest whose conformation is unavailable, said method comprising introducing amino acid changes into said protein to create an engineered protein, said engineered protein having the conformation of the native protein wherein said conformation of the engineered protein is confirmed by binding with an interacting molecule which binds with the native protein.
2. The method of Claim 1, wherein said interacting molecule is an antibody.
3. The method of Claim 2, wherein said antibody is a monoclonal antibody.
4. The method of Claim 1, wherein said amino acid changes are made to increase levels of essential amino acids in the engineered protein.
5. The method of Claim 4, wherein said essential amino acid is selected from the group consisting of methionine, tryptophan, lysine, valine, phenylalanine, isoleucine, leucine, threonine and cysteine.
6. The method of Claim 5, wherein said essential amino acid is methionine.
7. The method of Claim 1, wherein said amino acid changes are introduced into predetermined sites.
8. The method of Claim 7, wherein said predetermined site is determined by secondary structure prediction or homology comparison.
9. The method of Claim 1, wherein said amino acid changes are introduced at random.
10. The method of Claim 9, wherein said amino acid changes are produced by mutagenic PCR, DNA shuffling, or phage display methodology.
11. The method of Claim 10, wherein correctly folded variants are confirmed by filter lift assay or ELISA.

12. The method of Claim 4, wherein said essential amino acids are increased to represent 5% of the total amino acid content of the protein.
13. The method of Claim 4, wherein said essential amino acids are increased to represent 10% of the total amino acid content of the protein.
14. The method of Claim 1, wherein said protein is vegetative storage protein.
15. A method for altering amino acid composition of a native protein of interest, said method comprising introducing amino acid changes into said protein to increase nutritional value to create an engineered protein, said engineered protein to create an engineered protein, said engineered protein having the conformation of the native protein wherein said conformation of the engineered protein is confirmed by binding with an interacting molecule which binds with the native protein.
16. The method of Claim 15, wherein said interacting molecule is an antibody.
17. The method of Claim 16, wherein said antibody is a monoclonal antibody.
18. The method of Claim 15, wherein said amino acid changes are made to increase levels of essential amino acids in the engineered protein.
19. The method of Claim 18, wherein said essential amino acid is selected from the group consisting of methionine, tryptophan, lysine, valine, phenylalanine, isoleucine, leucine, threonine and cysteine.
20. The method of Claim 19, wherein said essential amino acid is methionine.
21. The method of Claim 15, wherein said amino acid changes are introduced into predetermined sites.
22. The method of Claim 21, wherein said predetermined site is determined by secondary structure prediction or homology comparison.

23. The method of Claim 15, wherein said amino acid changes are introduced at random.
24. The method of Claim 23, wherein said amino acid changes are produced by mutagenic PCR, DNA shuffling, or phage display methodology.
25. The method of Claim 24, wherein correctly folded variants are confirmed by filter lift assay or ELISA.
26. The method of Claim 18, wherein said essential amino acids are increased to represent 5% of the total amino acid content of the protein.
27. The method of Claim 18, wherein said essential amino acids are increased to represent 10% of the total amino acid content of the protein.
28. The method of Claim 15, wherein said protein is vegetative storage protein.
29. An engineered protein having altered amino acid composition, wherein said amino acid composition has been altered by introducing amino acid changes into said protein, wherein said engineered protein binds to an interacting molecule capable of binding with a corresponding native protein.
30. The protein of Claim 29 wherein said interacting molecule is an antibody.
31. The protein of Claim 30, wherein said antibody is a monoclonal antibody.
32. The protein of Claim 29, wherein said amino acid changes increase the levels of essential amino acids in the protein.
33. The protein of Claim 32, wherein said essential amino acid is selected from the group consisting of methionine, tryptophan, lysine, valine, phenylalanine, isoleucine, leucine, threonine and cysteine.
34. The protein of Claim 33, wherein said essential amino acid are increased to represent 5% of the total amino acid content of the protein.
35. The protein of Claim 32, wherein said essential amino acid are increased to represent 10% of the total amino acid content of the protein.

36. The protein of Claim 35, wherein said essential amino acid is methionine.
37. The protein of Claim 29, wherein said amino acid changes are introduced into predetermined sites.
38. The protein of Claim 37, wherein said predetermined site is determined by secondary structure prediction or homology comparison.
39. The protein of Claim 29, wherein said amino acid changes are introduced at random.
40. The protein of Claim 39, wherein said amino acid changes are produced by mutagenic PCR, DNA shuffling, or phage display methodology.
41. The protein of Claim 29, wherein said protein is vegetative storage protein.
42. A nucleotide sequence encoding the protein of Claim 29.
43. A nucleotide sequence encoding the protein of Claim 32.
44. A transformed plant containing within its genome the nucleotide sequence of Claim 42.
45. A transformed plant containing within its genome the nucleotide sequence of Claim 43.
46. A stably transformed plant having inserted into its genome a chimeric gene said gene encoding an engineered protein having altered amino acid composition wherein said protein, wherein said engineered protein binds to an interacting molecule which binds with a corresponding native protein.
47. The plant of Claim 46, wherein said amino acid changes increase the levels of essential amino acids in the protein.
48. The plant of Claim 47, wherein said essential amino acid is selected from the group consisting of methionine, tryptophan, lysine, valine, phenylalanine, isoleucine, leucine, threonine and cysteine.
49. The plant of Claim 48, wherein said essential amino acid are increased to represent 5% of the total amino acid content of the protein.

50. The plant of Claim 48, wherein said essential amino acid are increased to represent 10% of the total amino acid content of the protein.

51. The plant of Claim 46, wherein said plant is a dicot.

52. The plant of Claim 46, wherein said plant is a monocot.

53. The plant of Claim 52, wherein said monocot is maize.

54. The plant of Claim 52, wherein said dicot is soybean.

55. Seed of the plant of Claim 46.

56. Seed of the plant of Claim 52.

57. Seed of the plant of Claim 53.

VSP HOMOLOGIES

	1	5	10	15	20	25	30
VSP-b	RSSEV	KCAS	FR	LAVE	EA	HNIRAK	TIP
VSP-a	RTPEV	KCAS	WR	LAVE	EA	HNIRAK	TIP
T.phos		LKCTT	WR	FV	VE	TNNLS	PWK
Ph.vulg	SDTEV	RCAS	WR	LAVE	AQ	NI	FGFET
Ar.VSP		PNC	RSW	H	LG	FET	SNM
Ar.1A-1	S	I	N	Y	P	N	C
Ar17A-1	S	I	N	Y	A	N	C
	31	35	40	45	50	55	
VSP-b	EPI	KDY		INGE	FR	SD	SKIVNQQA
VSP-a	EAT	KEY		INGE	YR	SD	SKIVNQQA
T.phos	DYV	KEY		MVG	PGY	KMEI	DRVS
Ph.vulg	DAT	ANY		IEG	GQY	RSD	SKIVNQQA
Ar.VSP	AYV	EDY	LI	TS	KQY	QYD	SKIVNKEA
Ar.1A-1	AYV	EDY	LI	TS	KQY	QYD	SKIVNKEA
Ar17A-1	DYV	EDY	LI	TS	KQY	QYD	SKIVNKEA
	60	65	70	75	80	85	
VSP-b	REV		HHND	I	FI	FG	DNIVLSNIPYYEK
VSP-a	LEV		HPKD	TEV	FS	DNIVLSNIPYYEK	
T.phos	VDE	GDD	GR	EVW	F	DVDE	ILSNLPYYSDHR
Ph.vulg	RHV		HEND	VILE	N	DGTA	LSNIPYYSQHG
Ar.VSP	LAL	KND	TI	NVW	F	DLDD	ILSSIPYYAKYG
Ar.1A-1	LAL	KND	TI	NVW	F	DLDD	ILSSIPYYAKYG
Ar17A-1	LAL	KND	TI	NVW	F	DLDD	ILSSIPYYAKYG
	90	95	100	105	110	115	
VSP-b	YGVEE	FNET	LY	DEW	VN	KGD	APAL
VSP-a	YGVEE	FNET	LY	DEW	VN	KGN	APAL
T.phos	YGLEV	FDD	VE	F	DKW	VENGT	APAL
Ph.vulg	YGSEK	FDS	ERY	D	E	FVN	KGEAPAL
Ar.VSP	YGTENT	AAG	AY	WS	W	VSCE	TPGL
Ar.1A-1	YGTENT	AAG	AY	WS	W	VSCE	TPGL
Ar17A-1	YGTENT	AAG	AY	WS	W	VSCE	TPGL

TO FIG. 1A.

FIG. 1.

3 / 9


PROPOSED VSP β METHIONINE-ENRICHED VARIANTS

	1	5	10	15	20	25	30
VSP β	RSSEV	KCAS	FRLAVE	AHN	I	RAFKT	IPEECV
VSP β -Met10		M			M		M
VSP β -Met20		M			M		M
VSP β -Met30		M			M	M	M
	31	35	40	45	50	55	60
VSP β	EPTKDY	I	NGEQFR	SDSKT	DN	QQAFF	YASER
VSP β -Met10		M		M			M
VSP β -Met20	M	M		M			M
VSP β -Met30	M	M		M	M		M
	61	65	70	75	80	85	90
VSP β	EVHHND	I	FIFG	IDNT	VLSN	I	PYYEKHGYGV
VSP β -Met10	M	M	M	M			
VSP β -Met20	M	M	M	M			M
VSP β -Met30	M	M	M	M	M		M
	91	95	100	105	110	115	120
VSP β	EEFN	ETLY	DEWV	NKGD	APAL	PETL	KNYNKL
VSP β -Met10							
VSP β -Met20		M	M				
VSP β -Met30	M	M	M				
	121	125	130	135	140	145	150
VSP β	LSLGFK	I	VFL	SGRYL	DKMAV	TEANL	KKAGF
VSP β -Met10	M	M			M		M
VSP β -Met20	M	M	M	M	M	M	M
VSP β -Met30	M	M	M	M	M	M	M
	151	155	160	165	170	175	180
VSP β	HTWEQ	LILK	DPHL	IITPNA	L	SYKSAM	RENLL
VSP β -Met10						M	M
VSP β -Met20				M	M	M	M
VSP β -Met30			M	MM	M	M	M
	181	185	190	195	200	205	210
VSP β	RQGYR	I	VGII	IGDQW	SDL	LGDR	GESRTFKL
VSP β -Met10		M				M	
VSP β -Met20		M			M	M	M
VSP β -Met30		M			M	M	M
	211	215	218				
VSP β	PNP	MYI	IE				
VSP β -Met10	M	M					
VSP β -Met20	M	M					
VSP β -Met30	M	M					

FIG. 2.

SUBSTITUTE SHEET (RULE 26)


HYDROPATHY INDEX COMPUTATION FOR SEQUENCE VSPB.
TOTAL NUMBER OF AMINO ACIDS IS: 218.



HYDROPATHIC INDEX OF VSPB FROM AMINO ACID 1 TO AMINO ACID 218.
COMPUTED USING AN INTERVAL OF 9 AMINO ACIDS. (GRAVY=-4.95).

FIG. 3A.


HYDRPATHY INDEX COMPUTATION FOR SEQUENCE VSPM10.
TOTAL NUMBER OF AMINO ACIDS IS: 218.



HYDRPATHIC INDEX OF VSPM1 FROM AMINO ACID 1 TO AMINO ACID 218.
COMPUTED USING AN INTERVAL OF 9 AMINO ACIDS. (GRAVY=-5.52).

FIG. 3B.


HYDROPATHY INDEX COMPUTATION FOR SEQUENCE VSPM20.
TOTAL NUMBER OF AMINO ACIDS IS: 218.



HYDROPATHIC INDEX OF VSPM20 FROM AMINO ACID 1 AMINO ACID 210.
COMPUTED USING AN INTERVAL OF 9 AMINO ACIDS. (GRAVY=-5.68).

FIG. 3C.

HYDROPATHY INDEX COMPUTATION FOR SEQUENCE VSPM30.
TOTAL NUMBER OF AMINO ACIDS IS: 218.



HYDROPATHIC INDEX OF VSPM30 FROM AMINO ACID 1 TO AMINO ACID 218.
COMPUTED USING AN INTERVAL OF 9 AMINO ACIDS. (GRAVY=-5.31).

FIG. 3D.

Figure 4

VSP β -met10 sequence

*Sfi*I

1 GGCCCAGCCGGCCAGATCTTCGGAGATGAAATGCGCTAGCTTTAGGCTTGCTGTGGAAGC 60
CCGGGTCCGGCCGGTCTAGAAGCCTCTACTTTACGCGATCGAAATCCGAACGACACCTTCG

61 ACACAACATGCGAGCCTTTAAAACCATTCCTGAAGAGTGCATGGAACCAACAAAGGACTA 120
TGTGTTGTACGCTCGGAAATTTGGTAAGGACTTCTCACGTACCTTGGTTGTTTCCTGAT

121 CATGAATGGCGAACAAATTTCAATGGACTCTAAAACAGTTPACCAACAGGCCTTCTTTTA 180
GTACTTACCGCTTGTTAAAGCTTACCTGAGATTTTGTCAATTGGTTGTCCGGAAGAAAAT

181 TGCTAGTGAAATGGAAATGCATCACAACGACATGTTTATATTCGGCATGGATAACACCAT 240
ACGATCACTTTACCTTTACGTAGTGTTGCTGTACAAATATAAGCCGTACCTATTGTGGTA

241 GCTCTCTAATATCCCATACTATGAAAAACATGGATATGGGGTGGAGGAATTTAATGAAAC 300
CGAGAGATTATAGGGTATGATACTTTTTGTACCTATACCCACCTCCTTAAATTACTTTG

301 CTTATATGATGAATGGGTTAACAAGGGCGACGCACCGGCATTGCCAGAGACTCTTAAAAA 360
GAATATACTACTTACCCAATTGTTCCCGCTGCGTGGCCGTAACGGTCTCTGAGAATTTTT

361 TTACAACAAGCTGATGTCCCTTGGCTTCAAGATGGTATTCTTGTGAGGAAGGTACCTTGA 420
AATGTTGTTGCTGACTACAGGGAACCGAGTTCTACCATAAGAACAGTCCTTCCATGGAAC

421 CAAAATGGCCGTAAACAGAAGCAAACCTAATGAAGGCTGGCTTCCACACATGGGAGCAGTT 480
GTTTTACCAGCATTGTCTTCGTTTGGATTACTTCCGACCGAAGGTGTGTACCCTCGTCAA


481 AATTCTCAAGGATCCACATCTTATGACTCCAAATGCACCTTTCATACAAATCAGCAATGAG 540
TTAAGAGTTCCTAGGTGTAGAATACTGAGGTTTACGTGAAAGTATGTTTAGTCGTTACTC

541 AGAGAATATGTTGAGGCAGGGATACAGAATTGTTGGAATGATTGGTGATCAATGGAGCGA 600
TCTCTTATACAACCTCCGTCCCTATGTCTTAAACAACCTTACTAACCCTAGTTACCTCGCT

601 TCTGCTTGGAGACCACATGGGCGAATCTAGAACCTTTAAGCTTCCTAATCCCATGTACTA 660
AGACGAACCTCTGGTGTACCCGCTTAGATCTTGGAATTCGAAGGATTAGGGTACATGAT

661 CATGGAGGCGGCCGC 675
GTACCTCCGCCGGCG

NotI

**FIG. 5.**